

key

# MATH 135-08, 135-09 Calculus 1, Fall 2017

## Inverse Functions: Worksheet for Section 1.5

### The Inverse

One of the simplest ways to obtain a new function from an old one is to simply flip the domain and range. So if  $f(2) = 7$ , then the new function  $f^{-1}$ , called the **inverse of  $f$** , has  $f^{-1}(7) = 2$ . The function  $f^{-1}$  simply maps each element in the range of  $f$  back to the element in the domain it came from; it "inverts"  $f$ .

However, there will be a problem if an element  $b$  in the range has two or more elements in the domain, say  $a_1$  and  $a_2$ , that map to it (i.e.,  $f(a_1) = f(a_2) = b$ ). Then, the inverse of  $b$  would not be unique (it could be either  $a_1$  or  $a_2$ ), and  $f^{-1}$  would not be a function. To rectify this, we must assume that  $f$  is **one-to-one**, that is, each element in the range of  $f$  has one and only one pre-image that was sent to it in the domain.

For example, the function  $f(x) = x^2$  is not one-to-one because both  $-2$  and  $2$  are each sent to the same element in the range,  $f(-2) = f(2) = 4$ . This function fails the **horizontal line test** and is really two-to-one. However, if we restrict the domain to  $x \geq 0$  (think of erasing the left half of the parabola), then  $f$  becomes one-to-one and now the inverse is actually a function. You know it already as  $f^{-1}(x) = \sqrt{x}$ . By definition (i.e., restricting the domain of  $x^2$ ),  $\sqrt{x}$  only spits out non-negative values.



**Key Point:** The only functions with well-defined inverses are those that are **one-to-one**. They must pass the **horizontal line test**.

**Exercise 0.1** Which of the following functions are one-to-one on their full domains? Draw a few examples to illustrate the difference between a one-to-one function and a function that is not one-to-one.

a.  $f(x) = \frac{2}{3}x - 7$     b.  $F(x) = \frac{2}{3}$     c.  $g(x) = (x - 3)^2 + 5$      d.  $G(x) = (x - 3)^3 + 5$

e.  $h(\theta) = 2\sin(3\theta)$     f.  $H(\beta) = -4\tan(\beta)$     g.  $i(t) = |t + 1| - 4$      h.  $Y(t) = \sqrt{t + 1} - 4$

All other choices fail the horizontal line test.



Note that the notation for the inverse of  $f$  is not the usual exponent notation. In other words,

$$f^{-1} \neq \frac{1}{f}$$

The choice of  $-1$  as the exponent is mathematical shorthand for **inverse**. Don't confuse this!

Based on the definition of the inverse of a function, the following formulas should make sense:

$$\boxed{f^{-1}(f(x)) = x \quad \text{and} \quad f(f^{-1}(x)) = x.} \quad (1)$$

Simply put, the inverse of  $f$  reverses what  $f$  does to  $x$ . Likewise,  $f$  reverses the action of  $f^{-1}$ , that is, the inverse of  $f^{-1}$  is just  $f$ . Note that the domain of  $f^{-1}$  equals the range of  $f$ , and vice-versa.

Graphically, if  $(x, y)$  is a point on the graph of  $f$ , then  $(y, x)$  is a point on the graph of  $f^{-1}$ . Thus, to obtain the graph of  $f^{-1}$  from the graph of  $f$  (or vice versa), just reflect the graph of  $f$  about the line  $y = x$ . This is also how one obtains an analytic formula for the inverse of a function: interchange the variables  $x$  and  $y$  and solve for  $y$ .

**Exercise 0.2** Find the inverse of the function  $f(x) = 1 + \sqrt{2+3x}$ . What should the domain of  $f^{-1}$  be in order to have  $f$  as its inverse?

$y = 1 + \sqrt{2+3x}$ , Switch  $x$  and  $y$  and solve for  $y$ .

$$x = 1 + \sqrt{2+3y} \Rightarrow x-1 = \sqrt{2+3y} \Rightarrow 2+3y = (x-1)^2$$

$$\Rightarrow 3y = (x-1)^2 - 2 \Rightarrow y = \frac{(x-1)^2 - 2}{3} = \frac{1}{3}(x-1)^2 - \frac{2}{3}$$

$f^{-1}(x) = \frac{1}{3}(x-1)^2 - \frac{2}{3}$ . Domain of  $f^{-1} = \text{Range of } f$   
 $= [1, \infty)$   
 (draw graph, shift correctly)

### Inverse Trig Functions

The key to defining the inverse trig functions is to restrict the domains of the original trig functions in order to ensure that they are one-to-one. For example, the sine function is one-to-one on the domain  $-\pi/2 \leq \theta \leq \pi/2$  (check the graph). By making this restriction, we then **define** the range of the inverse sine function (also called the **arcsine function**) to be  $[-\pi/2, \pi/2]$ . The domain of the inverse sine function is  $[-1, 1]$  because this is precisely the range of the sine function.

**Key Point:** The inverse sine function, denoted  $\sin^{-1}(x)$ , inputs numbers between  $-1$  and  $1$  and outputs angles between  $-\pi/2$  and  $\pi/2$ . Thus,  $\theta = \sin^{-1}(x)$  if and only if  $\sin(\theta) = x$ . (Go backwards!)

The domains and ranges of the other inverse trig functions are given below:

- $\cos^{-1}(x)$ : Domain:  $[-1, 1]$  Range:  $[0, \pi]$
- $\tan^{-1}(x)$ : Domain:  $(-\infty, \infty)$  Range:  $(-\pi/2, \pi/2)$
- $\cot^{-1}(x)$ : Domain:  $(-\infty, \infty)$  Range:  $(0, \pi)$
- $\sec^{-1}(x)$ : Domain:  $(-\infty, -1] \cup [1, \infty)$  Range:  $[0, \pi/2) \cup (\pi/2, \pi]$
- $\csc^{-1}(x)$ : Domain:  $(-\infty, -1] \cup [1, \infty)$  Range:  $[-\pi/2, 0) \cup (0, \pi/2]$



**Exercise 0.3** Evaluate each of the following without using a calculator:

a.  $\sin^{-1}(1/\sqrt{2}) = \underline{\pi/4}$       b.  $\sin^{-1}(1) = \underline{\pi/2}$       c.  $\cos^{-1}(-\sqrt{3}/2) = \underline{5\pi/6}$   
 d.  $\tan^{-1}(-1) = \underline{-\pi/4}$       e.  $\sec^{-1}(2) = \underline{\pi/3}$       f.  $\csc^{-1}(-1) = \underline{-\pi/2}$

c. Solve  $\cos \theta = -\sqrt{3}/2$  with  $0 \leq \theta \leq \pi$ . Unit circle  $\Rightarrow$

f. Solve  $\csc \theta = -1$  or  $\frac{1}{\sin \theta} = -1$  with  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ ,  $\theta = 5\pi/6$ .  
 $\theta \neq 0$ .  
 $\Rightarrow \sin \theta = -1 \Rightarrow \theta = -\pi/2$

**Exercise 0.4** Explain why  $\pi/2$  must be excluded from the range of  $\sec^{-1}(x)$ . In other words, why does the equation  $\sec^{-1}(x) = \pi/2$  have no solution?

If  $\sec^{-1}(x) = \pi/2$ , then  $\sec \frac{\pi}{2} = x$ .

But  $\sec \frac{\pi}{2} = \frac{1}{\cos \frac{\pi}{2}} = \frac{1}{0} = \text{undefined}$ .

Thus, no solution exists.

**Exercise 0.5** Explain why  $\cos^{-1}(\cos(17\pi)) = \pi$  and not  $17\pi$ . What is  $\sin^{-1}(\sin(11\pi/3))$ ?

$\cos^{-1}(\cos 17\pi) = \pi$  because the output of  $\cos^{-1} x$  is always between 0 and  $\pi$ .  $\sin^{-1}(\sin \frac{11\pi}{3}) = -\pi/3$  because

$-\pi/2$  and  $11\pi/3$  are the same angle on the unit circle

and

$$-\frac{\pi}{2} \leq -\frac{\pi}{3} \leq \frac{\pi}{2}$$




Figure 1: The graph of the inverse tangent function  $y = \tan^{-1} x$  is cool. It maps the entire real line one-to-one and onto the open interval  $(-\pi/2, \pi/2)$ . Note the horizontal asymptotes at  $y = -\pi/2$  and  $y = \pi/2$ .

